

Review Article

Open Access, Volume 3

Complete Rupture of the Insertional Tendon of Biceps Brachii: What we Know

Nugent Luke*

Royal Shrewsbury Hospital, The Shrewsbury and Telford Hospital NHS Trust, Shrewsbury, SY3 8XQ, England.

Abstract

Purpose and objective: Ruptures of the distal biceps tendon present surgeons with a complex set of potential management options, and significant variation in practice exists. This review aims to succinctly collate the current evidence with regards to epidemiology, diagnosis, and management, in order to allow clinicians to inform themselves as to how best to deal with this increasingly frequently encountered phenomenon.

Background and principle results: This review focuses on an area of orthopaedics which demonstrates significant heterogeneity in its management. It was felt that the overwhelming amount of published work on the subject of distal biceps tendon ruptures had yet to be collated into a succinct article, and so literature review and summarisation was performed and presented herein.

Summary and major conclusions: The relative frequency of the injury, relevant functional anatomy, pathophysiology, management options, and notes on post-operative rehabilitation are included.

Keywords: Distal biceps; Tendon rupture; Biceps repair; Upper limb; Sports injuries; Biceps brachii.

Review

Functional anatomy of the distal insertion

The role of the biceps brachii (biceps) as a powerful supinator of the forearm has been well described [1] and will be self-evident to most. Despite sharing a 'common' distal tendinous insertion, the fibres of the long and short heads remain distinct, with the long head occupying a much larger ovoid footprint on the radial tuberosity, and the short inserting in a slender distally orientated outcropping [2]. The long head therefore influencing supination more so than the short head. In addition to the true bony attachment, the insertional tendon also gives of the lacertus

fibrosus. Also referred to as the bicipital aponeurosis, this thick fibrous structure blends with the deep fascia of the forearm and prevents over-lateralisation of the ulna during supination [3]. Its historical moniker of the *Grace à Dieu* fascia derives from the fact that it would protect the neurovascular structures of the antecubital fossa from inadvertent injury during supposedly therapeutic phlebotomy of the median cubital vein [4].

In ruptures of the distal biceps tendon distal to the lacertus, it acts like a vinculum, tethering the muscle and impeding complete retraction towards the shoulder. This accounts for the inconsistent appearance of the "reverse Popeye sign". It may also lead

Manuscript Information: Received: Apr 24, 2023; Accepted: May 17, 2023; Published: May 26, 2023

Correspondance: Nugent Luke, Royal Shrewsbury Hospital, The Shrewsbury and Telford Hospital NHS Trust, Shrewsbury, SY3 8XQ, England. Email: luke.nugent@doctors.org.uk

Citation: Luke N. Complete Rupture of the Insertional Tendon of Biceps Brachii: What we Know. *J Surgery*. 2022; 3(1): 1103.

Copyright: © Luke N 2023. Content published in the journal follows creative common attribution license.

to a false-negative “hook test”; the examiner may believe they can feel an intact insertional tendon when they are in fact palpating the remnant lacertus. In addition to these clinical findings the patient will present with pain and describes a “popping” or “tearing” sensation in the antecubital fossa. Radiological investigations are usually not required, but where doubt exists, ultrasound examination or magnetic resonance imaging are both equally as diagnostically useful (particularly in the case of incomplete rupture) [5]. Plain films are not usually helpful without the context of concomitant trauma, but may demonstrate hypertrophy about the radial tuberosity in the case of very longstanding insertional tendonopathy.

Seiler et al [6] described three zones for the insertional tendon of biceps, numbered from proximal to distal, and defined by their vascular supply; Zone 1 about the musculotendinous junction receiving its supply in much the same way as the muscular bulk proximal to it. I.e. via anastomosing arterioles which in turn each derive from the brachial artery itself. Zone 3 about the insertion’s footprint on the tuberosity is fed by the same branches of the posterior interosseous artery which supply the radial neck. The interposed Zone 2 is somewhat of a watershed area between the two supplies.

Tears; epidemiology and pathophysiology

Tears of the distal biceps tendon occur relatively infrequently, occurring in 0.8-1.9 patients in a population of 100,000 per year [7] (c.f tendo achillis ruptures; 21.5 per 100,000 per year [8], the most common tendon rupture) [9]. The typical patient is male, over 30 years old, and is predominantly affected on the dominant arm [10]. The Zone 2 vascular watershed is by far the most common rupture site. It is unfortunate that this particular region also appears to be most affected by mechanical impingement during pronosupination, resulting in fibrous attrition [6]. Factors thought to particularly increase risk are smoking and the use of corticosteroids [11]. Bilateral, or consecutive bilateral ruptures occur on occasion and are particularly associated with severe chronic kidney disease, hyperparathyroidism, and a recent history of quinolone-based antibiotic use [12]. Subjects who abuse testosterone and related compounds (anabolic steroids) for cosmetic purposes are also pre-disposed to tendon ruptures [13], possibly due to dysplastic organisation of collagen fibrils. Perhaps of most importance however is a pattern of acute injury on a background of overuse – weightlifters and manual labourers being particularly affected; An excessive eccentric loading force acts at the elbow as it is brought from extension into flexion [14]; the “flexed elbow unacceptably challenged”. In 1953 Chevallier [15] described a two-stage pathophysiological model which remains consistent with a contemporary understanding of the disease process. An interstitially weakened tendon ruptures acutely, and thereafter the lacertus fibrosis may also be torn as a result of un-opposed muscular contraction.

Conservative management and its outcomes

Many patients with an acute painful injury will be reluctant to “leave it alone” where an established surgical repair option exists. Outside of the context of patients with severe cognitive impairment or those whose medical comorbidities preclude operative intervention, conservative management of these injuries is generally reserved for the so-called “low-demand” patient [16]. Such

subjective evaluations of patient needs are inherently coloured by clinicians’ inherent biases; in reality there are very few people whose quality of life is not heavily dependent on their normal upper limb function. Counselling patients as to their options is also made difficult by the scarcity of evidence. No randomised clinical trials exist in the literature, and although some retrospective studies are reported, they are either underpowered [17,18], or effectively uncontrolled [19]. However what they can tell us is that outcomes for conservatively managed patients are generally poor [20], and although as much as 88% of elbow flexion power may be retained following complete distal biceps tendon rupture (compared with the uninjured arm), supination power and strength in resisted supination can be reduced to as little as 65% and 14% respectively [21]. Another point to consider is that conservatively managed patients who initially cope well may ultimately re-present with accelerated rotator cuff degeneration after adapting to their loss of supination strength by increasingly relying on external rotation at the shoulder as a compensatory technique [22].

Surgical management; History

Storhsin first identified the lesion at autopsy in 1842 [23]. Case reports on living subjects date back to the late 19th century [24], and by the 1950s about 100 cases had been described [25]. Even at that stage there is debate in the literature as to appropriate management techniques. Some cautioned that careful selection of operative candidates was imperative given the relatively minimal disability encountered by conservatively managed patients [26]. The earlier operative techniques consisted of either passing a heavy suture through the proximal part of the tear and attempting to approximate its anatomical footprint at the bicipital tuberosity by tying the suture ends about the whole of the proximal radius [27], or an alternative in the form of tendodesis of the distal biceps tendon to the brachialis (or to the ulna itself) [28]. This went some way to restoring strength in flexion but naturally could not be expected to assuage the supination deficit. Both techniques suffered from high failure rates, and disabling neurovascular complications were not unusual [29].

Modern surgical management

Surgical repair aiming to restore anatomy and function now represents the treatment goal for the majority of patients. Some technical variation exists in how surgeons go about achieving this, namely in the approach to fixation, as well as in the materials utilised.

One- versus two-incision techniques

Retrieval of the tendon stump and its re-attachment to the radius can be achieved by various approaches, but broadly speaking either one or two incisions are made. In the single incision technique, this may take the form of a small transverse incision just distal to the antecubital fossa, or a more extensile S-shaped incision can be used in cases where it is felt the stump may have more markedly retracted up the arm [30]. This technique is associated with a lower rate of heterotopic ossification and radio-ulnar synostosis (which can necessitate re-operation) than is seen in the dual incision alternative [31].

In the two incision variant, a second site is created over the radial head in order to receive material passed through a bony tunnel made through the proximal radius. This was developed with the

intention of more closely approximating the native anatomy [32], but recent meta-analysis has confirmed it also carries a lower risk of neurovascular injury [33], particularly to the lateral cutaneous nerve of the forearm (the Lateral Antebrachial Cutaneous Nerve, LACN). The same analysis also found no significant difference between the two techniques in terms of restoration of supination strength. The single incision approach was found to have better results in terms of range of motion in flexion and pronation, but this was caveated by heterogeneity in the rehabilitation regimens employed in the various included studies. Fortunately when LACN injuries do occur, they frequently take the form of self-limiting traction neuropraxiae [34], however more serious and disabling nerve injuries are also encountered in both approaches.

Surgical management; Fixation technique

Various methods of re-attaching the tendon have been proposed and remain in use. For primary repair of the native tendon, it can either be anchored to its footprint with an interference screw, or breasted with suture material which itself is made fast to the radius by Suture Anchors (SA), trans-osseous sutures (TO) or via an Endo-Cortical Button (ECB). In the single incision approach, an ECB is sometimes passed through the proximal cortex only, with the button sitting within the intramedullary canal of the radius.

There is a significant body of literature comparing the efficacy of these various materials. In cadaveric biomechanical studies, ECB has been found to be stronger than TO [35] but no observable real-world clinical differences were found in a retrospective cohort study [36]. Another biomechanical study found no difference in failure rates for SA versus ECB [37], and once again no real-world outcome differences were noted in a clinical study [38]. It would seem reasonable to conclude that when it comes to materials selection for these cases, *the best way is the way you know best*.

Post-operative rehabilitation

The ultimate goal of both repair and rehab is to enable the patient to return to work and recreational activities as quickly and as safely as possible. The exact nature of the rehabilitation programme advised will vary by centre, surgeon, and repair technique employed. In general however, an initial period of immobilisation is employed to protect the wound. This is followed by limited passive movement, and extension at the elbow may be restricted by a lockable range-of-motion or elbow hinge brace. Thereafter strengthening can begin. Biomechanical analyses have demonstrated that pull-out type failure is unlikely during physiological biceps contraction for various repairs [39], and this may reassure surgeons who are reluctant to “let them go” in the immediate post-operative period, but restricting higher intensity activities such as weights training is naturally a sensible precaution. Physiotherapists have demonstrated good outcomes with sequential, criterion-based, progressive rehabilitation programmes [40].

It is difficult to determine whether patients can expect their outcome to reflect their pre-injured state. Although anecdotally some patients report a return to competitive body-building and other high-demand work, it may be advisable to manage patients' expectations to a certain extent, but to what degree will always be

a function of individual experience.

Conclusion

- Distal biceps tendon rupture is relatively uncommon.
- Clinical assessment is usually sufficient in order to establish a diagnosis.
- Where doubt exists, ultrasound examination or magnetic resonance imaging may be sought.
- It is highly disabling and most patients benefit from operative repair.
- No single repair material has demonstrated superiority.
- Single incision approaches result in less heterotopic ossification.
- Dual incision approaches have fewer neurovascular complications.
- Structured rehabilitation under a physiotherapist's supervision is vital.

Declarations

Acknowledgements: On this occasion, there is no-one I wish to acknowledge. Thank you.

Conflict of interest: There are no conflicts of interest.

Funding/sponsorship: This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

Informed consent (Patient/Guardian): n/a, no patient data used Institutional Ethical Committee Approval; n/a.

Authors contribution: Single author.

References

1. Kleiber T, Kunz L, Disselhorst-Klug C. Muscular coordination of biceps brachii and brachioradialis in elbow flexion with respect to hand position. *Frontiers in Physiology*. 2015; 6: 215.
2. Jarrett CD, Weir DM, Stuffman ES, Jain S, Schmidt CC. Anatomic and biomechanical analysis of the short and long head components of the distal biceps tendon. *J Shoulder Elbow Surg*. 2012; 21: 942-948.
3. Deopujari R, Quadir N, Athavale S, Gajbhiye V, Kotgirwar S. Variant Bicipital Aponeurosis: A Cadaveric Study. *PSJR*. 2014; 7: 43-46.
4. Ellis H. The antecubital fossa. *Surgery*. 2010; 28: E1-E9.
5. Ramsey ML. Distal biceps tendon injuries: diagnosis and management. *J Am Acad Orthop Surg*. 1999; 7: 199-207.
6. Seiler JG, Parker LM, Chamberland PD, Sherbourne GM, Carpenter WA. The distal biceps tendon. Two potential mechanisms involved in its rupture: arterial supply and mechanical impingement. *J Shoulder Elbow Surg*. 199; 4: 149-156.
7. Safran MR, Graham SM. Distal biceps tendon ruptures: incidence, demographics, and the effect of smoking. *Clin Orthop Relat Res*. 2002; 404: 275-283.
8. Lantto I, Heikkinen J, Flinkkilä T, Ohtonen P, Leppihähti J. Epide-

- miology of Achilles tendon ruptures: increasing incidence over a 33-year period. *Scand J Med Sci Sports* 2015; 25: e133-e138.
9. Park SH, Lee HS, Young KW, Seo SG. Treatment of Acute Achilles Tendon Rupture. *Clin Orthop Surg.* 2020; 12: 1-8.
 10. Kelly MP, Perkinson SG, Ablove RH, Tueting JL. Distal Biceps Tendon Ruptures: An Epidemiological Analysis Using a Large Population Database. *Am J Sports Med.* 2015; 43: 2012-2017.
 11. Thomas JR, Lawton JN. Biceps and Triceps Ruptures in Athletes. *Hand Clin.* 2017; 33: 35-46.
 12. Michna H. Tendon injuries induced by exercise and anabolic steroids in experimental mice. *Int Orthop.* 1987; 11: 157-162.
 13. Visuri T, Lindholm H. Bilateral distal biceptendon avulsions with use of anabolic steroids. *Med Sci Sports Exerc.* 1994; 26: 941-944.
 14. Miyamoto RG, Elser F, Millett PJ. Distal biceps tendon injuries. *J Bone Joint Surg Am.* 2010; 92: 2128-2138.
 15. Chevallier CH. Sur un cas de désinsertion du tendon bicipital inferieur. *Mm Acad de Chir.* 1953; 79: 137-139.
 16. Miyamoto RG, Elser F, Millett PJ. Distal biceps tendon injuries. *J Bone Joint Surg Am.* 2010; 92: 2128-2138.
 17. Baker BE, Bierwagen D. Rupture of the distal tendon of the biceps brachii. Operative versus non-operative treatment. *J Bone Joint Surg AM.* 1985; 67: 414-417.
 18. Chillemi C, Marinelli M, De Cupis V. Rupture of the distal biceps brachii tendon: conservative treatment versus anatomic reinsertion – clinical and radiological evaluation after 2 years. *Arch Orthop Traum Surg.* 2007; 127: 705-708.
 19. Freeman CR, McCormick KR, Mahoney D, Baratz M, Lubahn JD. Nonoperative treatment of distal biceps tendon ruptures compared with a historical control group. *J Bone Joint Surg Am.* 2009; 91: 2329-2334.
 20. Hetsroni I, Pilz-Burstein R, Nyska M, Back Z, Barchilon V, Mann G. Avulsion of the distal biceps brachii tendon in middle-aged population: is surgical repair advisable? A comparative study of 22 patients treated with either nonoperative management or early anatomical repair. *Injury.* 2008; 39: 753-760.
 21. Cerciecki S, Visonà E, Corona K, Filho PRR, Carbone S. The Treatment of Distal Biceps Ruptures: An Overview. *Joints.* 2018; 6: 228-231.
 22. Meherin JM, Kilgore ES. The treatment of ruptures of the distal biceps brachii tendon. *Am J Sur.* 1960; 99: 636-640.
 23. Gilcreest EL. Rupture of muscles and tendons, particularly subcutaneous rupture of the biceps flexor cubiti. *J Amer Med Assoc.* 1925; 84: 1819-1822.
 24. Johnson AB. Avulsion of biceps tendon from the radius. *N Y Med J.* 1897; 66: 261-262.
 25. Thomas WJJ, Avulsion of the distal tendon of biceps brachii from the radial tuberosity. *South African Medical Journal.* 1958; 32: 1040-1042.
 26. Waugh RL, Hathcock TA, Elliot JL. Ruptures of muscles and tendons. *Surgery.* 1949; 25: 370-392.
 27. Lee H. Traumatic avulsion of tendon of insertion of biceps brachii. *Am J Surg.* 1951; 82: 290-292.
 28. Kron SD, Satinsky VP. Avulsion of the distal biceps brachii tendon. *Am J Surg.* 1954; 88: 657-659.
 29. Platt H, Observations on some tendon ruptures. *BMJ.* 1931; 1: 611-615.
 30. Camp CL, Voleti PB, Corpus KT, Dines JS. Single-Incision Technique for Repair of Distal Biceps Tendon Avulsions With Intramedullary Cortical Button. *Arthrosc Tech.* 2016; 5: e303-e307.
 31. Amin NH, Volpi A, Lynch TS, Patel RM, Cerynik DL, Schickendantz MS, et al. Complications of Distal Biceps Tendon Repair. *Orthop J Sports Med.* 2016; 4: 2325967116668137.
 32. Boyd HB, Anderson LD. A method for reinsertion of the distal biceps brachii tendon. *J Bone Joint Surg Am.* 1961; 43: 1041-1043.
 33. Castione D, Mercurio M, Fanelli D, Cosentino O, Gasparini G, Galasso O. Single- versus double-incision technique for the treatment of distal biceps tendon rupture. A systematic review and meta-analysis of comparative studies. *BJJ.* 2020; 102-B: 1608-1617.
 34. Carroll MJ, Dacambra MP, Hildebrand KA. Neurologic Complications of Distal Biceps Tendon Repair With 1-Incision Endobutton Fixation. *Am J Orthop.* 2014; 43: e159-e162.
 35. Kettler M, Tingart MJ, Lunger J, Kuhn LV. Reattachment of the distal tendon of biceps. Factors affecting the failure strength of the repair. *BJJ.* 2008; 90-B: 103-106.
 36. Recordon JA, Misur PN, Isaksson F, Poon PC. Endobutton versus transosseous suture repair of distal biceps rupture using the two-incision technique: a comparison series. *J Shoulder Elbow Surg.* 2015; 24: 928-933.
 37. Spang JT, Weinhold PS, Karas SG. A biomechanical comparison of EndoButton versus suture anchor repair of distal biceps tendon injuries. *J Shoulder Elbow Surg.* 2006; 15: 509-514.
 38. Reichert P, Królikowska A, Kentel M, Witkowski J, Gnus J, Satora W, Czamara A. A comparative clinical and functional assessment of cortical button versus suture anchor in distal biceps brachii tendon repair. *J Orthop Sci.* 2019; 24: 103-108.
 39. Rose DM, Archibald JD, Sutter EG, Belkoff SM, Wilckens JH. Biomechanical analysis suggests early rehabilitation is possible after single-incision EndoButton distal biceps repair with FiberWire. *Knee Surgery, Sports Traumatology, Arthroscopy.* 2011; 19: 1019-1022.
 40. Logan CA, Shahien A, Haber D, Foster Z, Farrington A, Provencher MT. Rehabilitation following distal biceps repair. *Int J Sports Phys Ther.* 2019; 14: 308-317.